INAF ISTITUTO NAZIONALE DI ASTROFISICA COSMIC RAYS: SHAPING-CHEMISTRY IN STAR-FORMING REGIONS MARCO PADOVANI

Impact of Low-Energy Cosmic Rays on Star Formation


Marco Padovani¹ · Alexei V. Ivlev² · Daniele Galli¹ · Stella S.R. Offner³ · Nick Indriolo⁴ · Donna Rodgers-Lee⁵ · Alexandre Marcowith⁶ · Philipp Girichidis⁷ · Andrei M. Bykov⁸ · J.M. Diederik Kruijssen⁹

Received: 17 January 2020 / Accepted: 27 February 2020 © Springer Nature B.V. 2020

Abstract In recent years, exciting developments have taken place in the identification of the role of cosmic rays in star-forming environments. Observations from radio to infrared wavelengths and theoretical modelling have shown that low-energy cosmic rays (< 1 TeV) play a fundamental role in shaping the chemical richness of the interstellar medium, determining the dynamical evolution of molecular clouds. In this review we summarise in a coherent picture the main results obtained by observations and by theoretical models of propagation and generation of cosmic rays, from the smallest scales of protostars and circumstellar discs, to young stellar clusters, up to Galactic and extragalactic scales. We also discuss the new fields that will be explored in the near future thanks to new generation instruments, such as: CTA, for the γ -ray emission from high-mass protostars; SKA and precursors, for the synchrotron emission at different scales; and ELT/HIRES, JWST, and ARIEL, for the impact of cosmic rays on exoplanetary atmospheres and habitability.

Other reviews

- Grenier et al. (2015) The nine lives of cosmic rays in galaxies
- Gabici (2022) Low-energy cosmic rays: regulators of the dense interstellar medium
- Owen et al. (2023) Cosmic-ray processes in galactic ecosystems



CHAPTER

Cosmic rays: physics, chemistry, and computational challenges

8

Marco Padovani^a and Brandt Gaches^{b,c,d}

^aINAF Osservatorio Astrofisico di Arcetri, Firenze, Italy

^bI. Physikalisches Institut, Universität zu Köln, Köln, Germany

^cDepartment of Space, Earth and Environment, Chalmers University of Technology, Gothenburg, Sweden ^dCenter of Planetary Systems Habitability, The University of Texas at Austin, Austin, TX, United States

The interstellar medium is immersed in a bath of cosmic rays: energetic charged particles with a non-thermal velocity distribution that exist in rough equipartition with the thermal and magnetic energies. Observations have shown that cosmic rays follow power-law distributions in energy from 1 GeV to 10 EeV with a turnover around 1 GeV. Cosmic rays are thought to have a wide range of effects on the interstellar medium and galaxies, such as ionization and heating of gas, pressure which aids in lifting gas out of galaxies, producing turbulence through instabilities in magneto-hydrodynamic turbulence. In this chapter, we will focus on the first of these and refer the reader to the many excellent reviews of the later effects (e.g., Strong et al., 2007; Zweibel, 2013; Grenier et al., 2015; Schlickeiser, 2015). We also refer the reader to Schlickeiser (2002) and Longair (2011) for more thorough monographs on cosmic-ray transport and high-energy phenomenon.





FIGURE 8.1

Emission processes diagnostics of cosmic-ray astrochemistry, with key current and proposed/future telescopes denoted at their respective wavelength regimes.


Astrochemical Modeling. https://doi.org/10.1016/B978-0-32-391746-9.00016-X Copyright © 2024 Elsevier Inc. All rights reserved. 189

SOME STATISTICS (end of 2022) Observations Theory Laboratory Astrochemistry Simulations 15 papers / yr 1980S PAPERS RELATED TO **COSMIC RAYS AND** STAR FORMATION 19 papers / yr 1990S decade 30.3 papers / yr 2000S 76.6 papers / yr 2010S 125 500 625 1000 250 375 750 875 0 no. of papers


SOME KEY ASPECTS IN GALACTIC STAR FORMATION




COSMIC-RAY FLUX



COSMIC-RAY FLUX



$$(p,e) + H_2 \rightarrow H_2^+ + e'' + (p',e')$$



$$(p,e) + H_2 \rightarrow H + H + (p',e')$$


-16.0

-17.0

-18.0

-18.5

Padovani+ (2018b) \log_{10} [E/eV]





$$p + H_2 \rightarrow H_2^+ + H_{fast} \qquad (\sigma_{e.c.}^p)$$

$$H_{fast} + H_2 \rightarrow H_{fast} + H_2^+ + e \qquad (\sigma_{ion}^H)$$

$$\rightarrow p + H_2 + e \qquad (\sigma_{self-ion}^H)$$


$$\rightarrow H_{fast} + H + H \qquad (\sigma_{diss}^H)$$






- **CHEMISTRY**;
- **♣ DYNAMICS OF STAR FORMATION;**
- ***** HEATING;
- **♣ DUST GRAIN CHARGE;**
- **SYNCHROTRON EMISSION**;
- * ATOMIC HYDROGEN FRACTION;
- **SPALLATION**;
- **♣** GAMMA-RAY EMISSION;







- **♣ DYNAMICS OF STAR FORMATION**;
- ***** HEATING;
- **♣ DUST GRAIN CHARGE**;
- **SYNCHROTRON EMISSION**;
- * ATOMIC HYDROGEN FRACTION;
- **SPALLATION**;
- **♣** GAMMA-RAY EMISSION;






COSMIC-RAY SIGNATURES

how can we reassemble the cosmic-ray spectrum from observations?


we have information on different energy ranges of the cosmic-ray flux using different observational techniques



* for $B_{\perp} \approx 2 - 20 \ \mu G$ and $\nu \approx 0.1 - 10 \ GHz$



COSMIC RAYS: CHARGED PARTICLES OUT OF THE THERMAL DISTRIBUTION



Maxwell-Boltzmann distribution

$$j_{\rm M} = \frac{c n_{\rm H_2} x E \exp[-E/(k_{\rm B}T)]}{\sqrt{2m(\pi k_{\rm B}T)^3}}$$

$$[j_{\rm M}] = \frac{\text{particles}}{\text{energy} \times \text{area} \times \text{time} \times \text{solid angle}}$$


COSMIC RAYS: CHARGED PARTICLES OUT OF THE THERMAL DISTRIBUTION

$$j_{\rm M} = \frac{c n_{\rm H_2} x E \exp[-E/(k_{\rm B}T)]}{\sqrt{2m(\pi k_{\rm B}T)^3}}$$


$$n(\mathrm{H_2})=500~\mathrm{cm^{\text{-}3}}$$
 ζ OPH $x=10^{\text{-}4}$ $T=50~\mathrm{K}$


$$n(\mathrm{H_2})=10^4~\mathrm{cm}^{-3}$$
TMC-1 $x=10^{-7}$ $T=10~\mathrm{K}$


$$n(\mathrm{H_2}) = 10^4 \mathrm{~cm^{-3}}$$
 $J \mathrm{FT}$ $x = 0.5$
 $T = 10^4 \mathrm{~K}$




COSMIC RAY: CHARGED PARTICLE OUT OF THE THERMAL DISTRIBUTION









AUGUST, 25TH 2012: VOYAGER 1 CROSSED THE HELIOPAUSE







still... variation in the magnetic field direction not seen so far



WHAT VOYAGER SPACECRAFTS ARE TELLING US

The two spacecraft share comparable cosmic-ray fluxes (90% agreement) in spite of the different crossing points of the heliopause.







COSMIC-RAY IONISATION RATE

number of H₂ ionisation per unit time

key-brick parameter





COSMIC-RAY IONISATION TRACERS

diffuse clouds

OH, HD, NH

Black & Dalgarno (1977)

Hartquist+ (1978)

van Dishoeck & Black (1986)

Federman+ (1996)

H_3^+, OH^+, H_2O^+

Geballe+ (1999),

McCall+ (2003),

Indriolo+ (2009,2012,2015),

Neufeld+ (2010), Gerin+ (2010)

ArH⁺

Neufeld & Wolfire (2017)

CO, OH, CH, HCO⁺

Luo+ (2023a,b)

dense clouds

HCO^+, DCO^+

Guélin+ (1977), Caselli+ (1998)

Maret & Bergin (2007), Cabedo+ (2023)

Pineda+ (2024)

N₂H⁺, HCO⁺, N₂D⁺, DCO⁺

 oH_2D^{\dagger}

Bovino+ (2020)

Sabatini+ (2023)

Redaelli+ (2024)

Redaelli+ (2021)

sulfur chemistry

Fuente+ (2016)

PO⁺

Rivilla+ (2022b)

H_2

Bialy (2020)

Padovani+ (2022)

Bialy, Belli, Padovani (2022)

Bialy+ (2025)

circumstellar discs

 HCO^+, N_2H^+

 oH_2D^+

Ceccarelli+(2004)

Cleeves+ (2014)

protostellar jets (shocks)

 HCO^+, N_2H^+

Podio+ (2014),

Ceccarelli+ (2014)

Rab+ (2017)

HC_3N , HC_5N

Fontani+ (2017)

 $c-C_3H_2$


Favre+ (2018)

ESTIMATES OF ζ FROM OBSERVATIONS


Sabatini+ 2023

HM SFRs

 oH_2D^+



KERNEL DENSITY ESTIMATE OF THE AVAILABLE ζ VALUES




COSMIC-RAY PROPAGATION INSIDE A MOLECULAR CLOUD

$$j(E_0, N = 0) = j^{IS}(E_0) \longrightarrow j(E, N)$$

Continuous Slowing-Down Approximation

- 1. straight-line propagation;
- 2. ΔE≪E.

$$L(E) = -\frac{1}{n} \frac{dE}{dx} = -\frac{dE}{dN}$$




$$N = \int_{E}^{E_0} \frac{\mathrm{d}E}{L(E)} \qquad \frac{\mathrm{d}E}{L(E)} = \frac{\mathrm{d}E_0}{L(E_0)}$$


$$j(E, N) dE = j^{IS}(E_0) dE_0$$

$$j(E,N) = j^{\mathrm{IS}}(E_0) \frac{\mathrm{d}E}{\mathrm{d}E_0} = j^{\mathrm{IS}}(E_0) \frac{L(E_0)}{L(E)} \longrightarrow \zeta(N) \propto \int j(E,N)\sigma(E) \,\mathrm{d}E$$


ENERGY LOSS FUNCTION PROTON + H₂




ENERGY LOSS FUNCTION ELECTRON + H2




ESTIMATES OF \(\Comparing models and observations \)





COSMIC-RAY FLUX BASED ON VOYAGER DATA






ESTIMATES OF & COMPARING MODELS AND OBSERVATIONS





PARAMETERISATION OF THE COSMIC-RAY SPECTRUM





$$j_k^{\text{IS}}(E) = C \frac{E^{\alpha}}{(E + E_0)^{\beta}} \text{ eV}^{-1} \text{ s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$$

Species k	\boldsymbol{C}	E_0 [MeV]	α	β – α
e	2.1×10^{18}	710	-1.3	3.2
$p \pmod{\mathscr{L}}$	2.4×10^{15}	650	0.1	2.7
$p \pmod{\mathcal{H}}$	2.4×10^{15}	650	-0.8	2.7


Most of the parameter space is dominated by the ionisation of CR protons and by the excitation of secondary electrons. For this reason we consider a single parameterisation for primary CR electrons.


ESTIMATES OF ζ comparing models and observations





ESTIMATES OF ζ comparing models and observations





LIMITATIONS OF CURRENT METHODS TO ESTIMATE ζ

diffuse clouds

▶ absorption line studies of H_3^+ , OH^+ , H_2O^+ and ArH^+

dense clouds

 $^{\triangleright}$ HCO⁺, DCO⁺, CO, oH₂D⁺ and more and more complex species

downsides


- need to have an early-type star in the background;
- ζ obtained from this method is proportional to $n(H_2)$;
- uncertainties in the local x_e that in turn depends on details of the IS UV attenuation in the cloud;

downsides

- require comprehensive and updated reaction networks;
- main source of uncertainty comes from the formation and destruction rates of some species;
- poorly constrained amount of carbon and oxygen depletion on dust grains;

A NEW METHOD TO ESTIMATE ζ IN MOLECULAR DENSE CLOUDS

First introduced by Bialy (2020), refined in PM+ (2022), and applied in Bialy+ (2022)




$$e + H_2 \rightarrow H_2^* + e'$$

lower degree of uncertainty

neither chemical networks nor abundances of other secondary species are involved

A NEW METHOD TO ESTIMATE ζ IN MOLECULAR DENSE CLOUDS



most efficient mechanism

Transition	Upper level (v, J)	Lower level (v', J')	λ [μ m]
(1-0)O(2)	(1,0)	(0,2)	2.63
(1-0)Q(2)	(1,2)	(0,2)	2.41
(1-0)S(0)	(1,2)	(0,0)	2.22
(1-0)O(4)	(1,2)	(0,4)	3.00

$$I_{ul}^{\text{dir}}(N) = \alpha_{ul} \frac{E_{ul}}{4\pi} \int_{0}^{N} \zeta_{\text{exc},u}(N') e^{-\tau_{d}(N')} x_{\text{H}_{2}}(N') dN'$$

 $\alpha_{ul} = A_{ul}/\Sigma_l A_{ul}$: probability to decay to state l given state u is excited

 E_{ul} : transition energy

 $\tau_d = \sigma_d N$: optical depth for dust extinction


 $\sigma_d \approx 4.5 \times 10^{-23} \ \mathrm{cm}^2$ per H nucleus (averaged over 2-3 µm)

 $x_{\rm H_2} = n_{\rm H_2}/(n_{\rm H} + 2n_{\rm H_2})$: molecular hydrogen fraction

$$\zeta_{\text{exc},u}(N_{\text{H}_2}) = 2\pi\ell \int j_k(E, N_{\text{H}_2}) \sigma_{\text{exc},u}(E) \, dE$$

 $\ell=1$ for primary CRs and $\ell=2$ for secondary electrons

A NEW METHOD TO ESTIMATE ζ IN MOLECULAR DENSE CLOUDS




017); Skrutskie et al. (2006)

(1-0)S(0) line at 2.22 µm

Magellan Infrared Spectrograph (MMIRS), mounted on the Multiple Mirror Telescope (MMT)

Each field of view is 40' x 40' - the 7' slit is shown in red



THE JWST REVOLUTION


	MMIRS @MMT	X-shooter @VLT	NIRSpec @JWST
transitions	(1-0)S(0)	(1-0)S(0) — (1-0)Q(2)	(1-0)O(2) – (1-0)Q(2) (1-0)S(0) – (1-0)O(4)
slit size	7'	11"x0.4"	3.4' x 0.27"
obs. time (3σ)	3 h (upper limits)	8 h	1.25 h


adding up the signal of e.g. 50 shutters (each shutter has a size of ~ 0.53"x0.27") it is possible to obtain a few independent estimates of I_{ν} and then of ζ .


E.g.: for Barnard 68 (distance of 125 pc) it is possible to obtain about 10 independent estimates of ζ .


Testing the presence of a gradient of ζ as predicted by models of attenuation of the interstellar CR spectrum

THE JWST REVOLUTION













IN MOLECULAR CLOUDS INDUCED BY COSMIC RAYS






UV H₂ LUMINESCENCE IN MOLECULAR CLOUDS INDUCED BY COSMIC RAYS





UV H2 LUMINESCENCE IN MOLECULAR CLOUDS INDUCED BY COSMIC RAYS



Photorates (dissociation and ionisation)

$$k_s^{\text{ph}}(N) = 4\pi \int \frac{J_{\nu}(N)\sigma_s^{\text{ph}}(\nu)}{h\nu} d\nu$$

 $\sigma_{\!\scriptscriptstyle S}^{
m ph}$ from the Leiden Observatory Database

$$k_s^{\text{ph}}(N) = k_{0,s}^{\text{ph}} \zeta_{H_2}(N)$$

parameterisation as a function of:

- interstellar CR spectrum;
- H₂ ortho-to-para ratio;
- dust composition (R_V : slope of extinction at visible wavelenths);
- ISRF


UV H2 LUMINESCENCE IN MOLECULAR CLOUDS INDUCED BY COSMIC RAYS

$R_V = 3.1$	$k_{0,s}^{ m pd}$		$k_{0,s}^{ m pd}$			$k_{0,s}^{ m pd}$		
Species s	o:p=0:1	o:p=1:0	Species s	o:p=0:1	o:p=1:0	Species s	o:p=0:1	o:p=1:0
Al	_	_	Ca ⁺	_	_	NH ₂ CHO	1.292(3)	1.294(3)
AlH	1.641(1)	1.644(1)	C1	_	_	NH_3^2	6.266(2)	6.093(2)
C	_	_	Cr	_	_	NO	1.715(2)	1.673(2)
\mathbf{C}_2	1.368(2)	1.368(2)	Fe	_	_	NO ₂	6.022(2)	6.046(2)
C_2^2H	8.184(2)	9.006(2)	Н	_	_	Na ²	_	_
$C_2^2H^-$	_	_	H-	_	_	NaCl	5.030(1)	5.036(1)
$C_2^2H_2$	1.086(3)	8.684(2)	H_2	4.048(2)	3.912(2)	NaH	2.323(2)	2.368(2)
$C_2^2H_3^2$	1.567(2)	1.566(2)	H_2^{\mp}	3.238(2)	3.290(2)	Ni	_	_
$C_2^2H_4^3$	1.450(3)	1.450(3)	H ₂ CO	6.247(2)	6.054(2)	О	_	_
$C_2^2H_5^T$	2.278(2)	2.262(2)	$H_2^{2}CS$	9.744(2)	9.734(2)	O_2	3.077(2)	3.106(2)
$C_2^T H_5^T OH$	1.466(3)	1.479(3)	$H_2^{\circ}O$	4.715(2)	4.670(2)	$O_2^{\tilde{\tau}}$	1.380(1)	1.420(1)
C_2H_6	1.264(3)	1.284(3)	H_2O_2	3.846(2)	3.893(2)	O_3	5.534(2)	5.633(2)
C_3	2.642(3)	2.600(3)	H_2S	1.387(3)	1.407(3)	OCS	2.319(3)	2.260(3)
C_3H_3	2.762(1)	2.765(1)	$H_3^{\tilde{7}}$	2.295(-1)	1.746(-1)	OH	2.646(2)	2.681(2)
$C_3^3H_7^3OH$	2.434(3)	2.470(3)	HC ₃ H	3.659(2)	3.708(2)	OH ⁺	8.375	8.063
$C_4^{\prime}H^{\prime}$	_	_	$HC_3^{\circ}N$	2.909(3)	2.901(3)	P	_	_
C_6H^-	_	_	HCN	1.052(3)	1.069(3)	PH	3.450(2)	3.318(2)
CH	4.598(2)	4.468(2)	HCO	2.211(2)	2.110(2)	PH ⁺	5.632(1)	5.736(1)
CH ⁺	2.045(2)	1.927(2)	HCO ⁺	3.049	3.770	Rb	_	_
CH_2	1.398(2)	1.383(2)	НСООН	9.896(2)	1.006(3)	S	_	_
CH_2^{7}	7.252(1)	7.117(1)	HC1	1.090(3)	9.841(2)	$\mid S_2 \mid$	3.176(1)	3.168(1)
CH_3	9.999(1)	1.058(2)	HCl ⁺	6.558(1)	6.413(1)	SĦ	3.782(2)	3.720(2)
CH_3 CHO	8.908(2)	8.989(2)	HF	1.022(2)	1.031(2)	SH ⁺	4.309(2)	4.220(2)
CH ₃ CN	1.955(3)	2.006(3)	HNC	9.632(2)	9.729(2)	SO	2.791(3)	2.827(3)
$CH_3^{\dagger}NH_2$	1.777(2)	1.755(2)	HNCO	1.431(3)	1.449(3)	SO_2	1.528(3)	1.534(3)
$CH_3^{\dagger}OCH_3$	9.651(2)	9.637(2)	HO_2	8.162(1)	8.122(1)	Si	_	_
CH ₃ OCHO	1.171(3)	1.177(3)	K	_	_	SiH	1.852(2)	1.867(2)
CH ₃ OH	8.862(2)	8.892(2)	Li	_	_	SiH ⁺	6.804(2)	6.981(2)
CH ₃ SH	1.197(3)	1.203(3)	LiH	1.522(2)	1.518(2)	SiO	5.673(2)	5.855(2)
CH_4	9.917(2)	9.972(2)	Mg	_	_	Ti	_	_
CH_4^+	1.852(2)	1.840(2)	MgH	8.085(1)	9.088(1)	Zn	_	_
CN	4.287(2)	4.350(2)	Mn	_	_	$c-C_3H$	1.668(2)	1.645(2)
CO	1.192(2)	6.450(1)	N	_	_	$c-C_3H_2$	2.523(2)	2.554(2)
CO ⁺	4.960(1)	5.239(1)	N_2	3.372(1)	4.319(1)	$1-C_3H$	1.326(3)	1.281(3)
CO_2	5.850(2)	6.443(2)	N_2O	9.807(2)	9.976(2)	$1-C_3H_2$	1.454(3)	1.432(3)
CS	6.416(2)	5.137(2)	NH	1.932(2)	1.860(2)	1-C ₄	4.637(2)	4.654(2)
CS_2	2.868(3)	2.687(3)	NH ⁺	1.678(1)	1.647(1)	$1-C_4H$	2.321(3)	2.314(3)
Ca	_	_	NH ₂	3.539(2)	3.559(2)	1-C ₅ H	4.787(1)	4.790(1)

$R_V = 3.1$	$k_{0,s}^{ m pi}$		_	$k_{0,s}^{ m pi}$		_	$k_{0,s}^{ m pi}$	
Species s	o:p=0:1	o:p=1:0	Species s	o:p=0:1	o:p=1:0	Species s	o:p=0:1	o:p=1:0
Al	1.023(3)	9.708(2)	Ca ⁺	1.585	1.465	NH ₂ CHO	3.739(2)	3.681(2)
AlH	6.841(1)	6.918(1)	C1	5.409(1)	5.322(1)	NH ₃	1.887(2)	1.896(2)
C	2.333(2)	2.381(2)	Cr	4.909(2)	4.967(2)	NO	1.793(2)	1.791(2)
C_2	2.193(2)	2.055(2)	Fe	2.064(2)	2.093(2)	NO ₂	1.032(2)	1.009(2)
${f C}_2 \\ {f C}_2 {f H}$	4.367(2)	4.197(2)	Н	4.078	3.615	Na ²	6.693	6.704
$C_2^2H^-$	6.119(2)	6.107(2)	H-	9.240(2)	9.247(2)	NaCl	_	_
$C_2^2H_2$	3.733(2)	3.603(2)	H_2	5.451(-2)	5.717(-2)	NaH	_	_
$C_2^2H_3^2$	1.335(3)	1.349(3)	H_2^{4}	_ ` ´		Ni	4.789(1)	4.887(1)
$C_2^2H_4^3$	2.668(2)	2.617(2)	H ₂ CO	2.703(2)	2.683(2)	О	2.668	2.358
$C_2^2H_5^4$	2.401(2)	2.425(2)	H ₂ CS	1.066(3)	1.065(3)	O_2	3.338(1)	2.656(1)
$C_2^2H_5OH$	3.620(2)	3.484(2)	H ₂ O	2.289(1)	2.106(1)	O_2^{\ddagger}	_ ` ´	_ ` ´
$C_2^2H_6^3$	1.827(2)	1.687(2)	$H_2^2O_2$	1.804(2)	1.730(2)	O_3^2	3.324(1)	3.048(1)
C_2^2	9.367(1)	9.051(1)	$H_2^2S^2$	5.771(2)	5.735(2)	OCS	5.284(2)	5.053(2)
$ C_3^2 $ $ C_3H_3$	1.022(3)	1.035(3)	$\begin{array}{c} H_2^2S \\ H_3^+ \end{array}$	_ ` ´	_ ` ´	ОН	1.662(1)	1.569(1)
$C_3^3H_7^3OH$	5.667(2)	5.468(2)	HC ₃ H	1.290(3)	1.306(3)	OH ⁺	_ ` `	_ ` `
$C_4^3H^{\prime}$	5.022(2)	5.013(2)	HC_3^3N	1.696(2)	1.575(2)	P	1.330(3)	1.359(3)
$C_6^4H^-$	3.042(2)	3.036(2)	HCN	1.401(1)	1.136(1)	PH	_ ` `	_
CH	1.621(2)	1.624(2)	HCO	5.843(2)	5.904(2)	PH ⁺	_	_
CH ⁺	_ ` ´	_ ` ´	HCO ⁺	_ ` `	_	Rb	1.095(1)	1.102(1)
CH_2	_	_	НСООН	2.035(2)	1.966(2)	S	7.989(2)	7.278(2)
CH_2^{\sharp}	_	_	HC1	4.771(1)	4.243(1)	S_2	1.667(3)	1.699(3)
CH_3^2	2.433(2)	2.458(2)	HCl ⁺			SH	9.110(2)	9.244(2)
CH_3 CHO	6.120(2)	6.256(2)	HF	1.026(-2)	9.336(-3)	SH ⁺	_ ` `	_ ` '
CH_3^3CN	9.682(1)	8.834(1)	HNC	1.518(2)	1.370(2)	SO	4.034(2)	4.298(2)
$CH_3^3NH_2$	1.157(3)	1.160(3)	HNCO	7.642(1)	7.172(1)	SO ₂	1.056(2)	9.662(1)
CH ₃ OCH ₃	8.670(2)	8.654(2)	HO ₂	5.597(1)	5.442(1)	Si ²	2.030(3)	2.000(3)
CH³OCHO	2.178(2)	2.114(2)	K	1.608(1)	1.606(1)	SiH	3.141(3)	3.146(3)
CH ₃ OH	2.314(2)	2.256(2)	Li	1.130(2)	1.126(2)	SiH ⁺		_
CH_3^3SH	1.354(3)	1.379(3)	LiH		_ ` `	SiO	_	_
CH_4^3	2.330(1)	2.039(1)	Mg	4.863(1)	4.797(1)	Ti	1.299(2)	1.312(2)
$CH_4^{\frac{7}{4}}$	_ ` `		MgH	_	_	Zn	4.198(2)	3.045(2)
CN ⁴	7.371	5.584	Mn	1.444(1)	1.453(1)	c-C ₃ H	9.757(2)	9.857(2)
CO	1.438(1)	1.029(1)	N	9.865(-1)	8.447(-1)	$c-C_3H_2$	1.227(3)	1.243(3)
CO ⁺	_	_	N_2	1.759(-1)	1.312(-1)	1-C ₃ H	9.998(2)	1.012(3)
CO_2	1.144(1)	9.550	N_2^2 O	1.079(2)	1.061(2)	1-C ₃ H ₂	_	_
CS ²	6.684(1)	6.861(1)	NH	7.819	6.735	$\begin{array}{ c c } 1-C_4 & & \\ \hline \end{array}$	_	_
CS ₂	2.770(2)	2.782(2)	NH ⁺	_	_	$1-C_4$ H	_	_
Ca ²	1.116(2)	1.110(2)	NH ₂	6.265(1)	6.122(1)	1-C ₅ H	_	_
			1 - 1 - 2	0.200(1)	(1 <i>)</i>	1 2511		

ESTIMATES OF & COMPARING MODELS AND OBSERVATIONS


this is *not*the end of the tale...




ESTIMATES OF ζ comparing models and observations

Values of ζ above the $\alpha=-1.2$ model cannot be explained by GCRs

an additional *local*CR source is needed



COSMIC-RAY IONISATION RATE INSIDE A MOLECULAR CLOUD




other sources of cosmic rays needed


COSMIC-RAY ACCELERATION SITES

Looking for shocks in/on: * accretion flows;


- protostellar surface;
- ▶ jet.


First-order Fermi acceleration (or Diffusive Shock Acceleration)







COSMIC-RAY ACCELERATION SITES













Other examples of synchrotron emission in protostellar jets \Rightarrow acceleration of local cosmic-ray electrons








Carrasco-González+ (2013) Rodríguez-Kamenetzky+ (2017)




Rodríguez-Kamenetzky+ (2016)

For high-mass protostellar jets see e.g. Araudo+ (2007), Bosch-Ramon+ (2010), Munar-Adrover+ (2011)

For a recent review, see Anglada+ (2018)







Carrasco-González+ (2013) Rodríguez-Kamenetzky+ (2017)

For high-mass protostellar jets see e.g. Araudo+ (2007), Bosch-Ramon+ (2010), Munar-Adrover+ (2011)

For a recent review, see Anglada+ (2018)

LOCALLY ACCELERATED COSMIC RAYS





$$U = 275^{+425}_{-50} \text{ km s}^{-1}$$


$$n/(10^6 \text{ cm}^{-3}) = 18.05^{+28.37}_{-13.67}$$


$$\tilde{P}/10^4 = 1.80^{+13.31}_{-1.10}$$

LOCALLY ACCELERATED COSMIC RAYS


fit to observations Meng+ (2019)


model Padovani+ (2019)





COSMIC-RAY ACCELERATION SITES






COSMIC-RAY ACCELERATION IN HII REGIONS

Chandra observations of the HII complex G5.89-0.39 and TeV γ -ray source HESS J1800-240B

- VLA 90 cm background red map
- Mopra NH₃(1,1) yellow contours
- ► HESS > 0.4 TeV white contours
- Chandra FOV green boxes

γ-rays as a TRIPLE
SIGNATURE of local
CRs?


10 out of 75 X-ray sources observed by Chandra could provide sufficient energy to account for a part of the GeV to TeV γ -ray emission in the source HESS J1800-240B. However, future arc-minute angular resolution γ -ray imaging will be needed to disentangle the potential γ -ray components powered by G5.89-0.39 from those powered by the W28 SNR.



in-situ CR acceleration already proposed by Gusdorf, Marcowith et al. (2015)

EFFECT OF STELLAR PARTICLES ON CIRCUMSTELLAR DISC CHEMISTRY

origin of the emission of HCO+(6-5) and $N_2H+(6-5)$



STELLAR PARTICLES: A NEW RESEARCH FIELD

